

Electric monopoles and the Montonen-Olive conjecture

M. Temple-Raston
Department of Mathematics,
Concordia University,
1455 de Maisonneuve Blvd. West,
Montreal, Quebec H3G 1M8 CANADA

February 1, 2008

Abstract

We introduce a topological field theory with a Bogomol'nyi structure permitting BPS electric, magnetic and dyonic monopoles. From the general arguments given by Montonen and Olive the particle spectrum and mass compare favourably with that of the intermediate vector bosons. In most, if not in all, of its essential features the topological field theory introduced here provides an example of a dual field theory, the existence of which was conjectured by Montonen and Olive.

In [5] Montonen and Olive conjectured the existence of a Lagrangian field theory ‘dual’ to Georgi-Glashow theory that would possess electric monopole solutions analogous to the BPS magnetic monopole. In favour of this conjecture Montonen and Olive argued that classical properties of the BPS monopole in the alleged dual field theory, namely the particle spectrum, the mass, and the long-range intermonopole force, correspond to the anticipated quantum properties of heavy gauge bosons (with a massless Higgs particle). The quantum corrections to the classical BPS monopole are unknown, but Montonen and Olive suggested that the corrections might be small, or, vanish entirely, due to symmetry. Lending some support to this expectation are recent studies of interacting classical BPS magnetic monopoles, showing

that a classical solitonic theory of BPS monopoles is capable of capturing dynamical behaviour usually associated with quantum mechanics [6].

Sixteen years after the conjecture was first made and despite the compelling evidence, not one explicit example of a dual field theory is known. This suggests that the dual field theory, if it exists, is structurally rather different from any theory previously considered. In [7] we argued that a topological field theory of a form first proposed by Horowitz [3] might lead to electric monopoles, and, perhaps, a dual field theory. The topological field theory was generalised in [8] to provide the Lagrangian with a Bogomol'nyi structure similar to the Bogomol'nyi structure leading to instantons in Yang-Mills theory. By virtue of the many similarities with self-dual instantons, solutions to the Bogomol'nyi equations in [8] are called 'topological instantons'. It is well-known that self-dual instantons and BPS magnetic monopoles are closely related by dimensional reduction with a gauge symmetry. By a similar process of dimensional reduction on topological instantons, a theory of topological monopoles is developed [9]. In this paper we study the theory of topological monopoles to see whether it can be viewed as a dual field theory.

Let G be a compact, semi-simple Lie group and P a principal G -bundle over a three-manifold M_3 . Denote the space of connections on P by $\mathcal{A}(P)$. We represent by E the vector bundle over M_3 associated to P by the adjoint representation. Associated with each connection $A \in \mathcal{A}(P)$ is an exterior covariant derivative D^A acting on sections of E . The covariant derivative defines a curvature H^A for the vector bundle E by $D^A D^A s = s H^A$, where s is a section of the vector bundle $E \rightarrow M$. The curvature H^A can be interpreted as a 2-form on M taking values in E . We also introduce an equivariant Lie algebra valued Higgs field, Φ_A , on M_3 associated with the connection A . Our theory is defined by the energy functional $\mathcal{E}(A, B, \Phi_A, \Phi_B)$ given by

$$\begin{aligned} \int_{M_3} & \langle (I_E \otimes K^B) \wedge (I_E \otimes D^B \Phi_B) \rangle - \langle (I_E \otimes K^B) \wedge (D^A \Phi_A \otimes I_E) \rangle \\ & - \langle (H^A \otimes I_E) \wedge (I_E \otimes D^B \Phi_B) \rangle + (A \leftrightarrow B, \Phi_A \leftrightarrow \Phi_B). \end{aligned} \quad (1)$$

I_E is the identity transformation on the adjoint bundle, E . Note that in (1) there are two curvatures H^A, K^B and two Higgs fields Φ_A, Φ_B corresponding to the two connections A, B . Finally, we assume that there is an invariant positive-definite inner product $\langle \rangle$ on E_x which varies continuously with $x \in M$. The last term in (1) symmetrises the energy functional in the dependent fields. The energy functional is rather complicated, but as we shall

see it is constructed so as to give certain geometric properties. The functional (1) can be derived from a four-dimensional topological field theory using dimensional reduction [9]. The three-dimensional theory can be examined independently of the four-dimensional theory, so a derivation of the three-dimensional theory is not needed here. The variational field equations arising from (1) are

$$\begin{aligned} D^B H^A &= 0, & D^B D^A \Phi_A &= [H^A, \Phi_B], \\ D^A K^B &= 0, & D^A D^B \Phi_B &= [K^B, \Phi_A]. \end{aligned} \quad (2)$$

The energy functional (1) can be rewritten as

$$\begin{aligned} \mathcal{E} = \int_{M_3} & \langle (H^A \otimes I_E - I_E \otimes K^B) \wedge (D^A \Phi_A \otimes I_E - I_E \otimes D^B \Phi_B) \rangle \\ & - \int_{M_3} \langle (D^A \Phi_A \otimes I_E) \wedge (H^A \otimes I_E) \rangle + (A \leftrightarrow B, \Phi_A \leftrightarrow \Phi_B). \end{aligned} \quad (3)$$

In this equation we begin to see the geometrical structure emerge out of the energy functional (1). Let E_A and E_B be the vector bundle E equipped with either the connection A or B , respectively. Recall that the curvature of the tensor product bundle $E_A \otimes E_B^*$ is given by $\Omega_{E_A \otimes E_B^*} = H^A \otimes I_E - I_E \otimes K^B$. This curvature expression appears in (3). By defining $\Phi = \Phi_A \otimes I_E - I_E \otimes \Phi_B$, then $D_{E_A \otimes E_B^*} \Phi = D^A \Phi_A \otimes I_E - I_E \otimes D^B \Phi_B$, and the first integral in (3) is therefore a topological invariant. The Bogomol'nyi equations arising from equation (1) are

$$\begin{aligned} H^A \otimes I_E &= I_E \otimes K^B \\ D^A \Phi_A \otimes I_E &= I_E \otimes D^B \Phi_B \end{aligned} \quad (4)$$

The first equation in (4) is therefore a zero curvature condition on the tensor product bundle $E_A \otimes E_B^*$. From an indices computation we conclude that

$$\begin{aligned} H^A &= K^B = iF I_E, \\ D^A \Phi_A &= D^B \Phi_B = iE I_E. \end{aligned} \quad (5)$$

F and E are a real-valued two-form and one-form on M_3 , respectively. Solutions to (5) automatically satisfy the variational field equations (2). Moreover, solutions to the first equation are projectively flat connections [4]. The projective flatness reduces the gauge group to $U(n)$. For line bundles this carries no extra information, but for bundles of rank greater than one projective flatness is a strong condition. Projectively flat connections are closely

related to Einstein-Hermitian connections, which in turn express in differential geometry the algebraic geometric concept of ‘stable vector bundle’ [4]. As a result, projectively flat connections have a well-behaved moduli space—like self-dual instantons and BPS magnetic monopoles. Unlike the theory of BPS magnetic monopoles, however, the energy functional (3) is saturated at the Bogomol’nyi energy with either Bogomol’nyi equation in (5) satisfied. Of course the field configurations must still satisfy the second-order variational field equations. The Bogomol’nyi energy is given by

$$\mathcal{E} = - \int_{M_3} \langle (D^A \Phi_A \otimes I_E) \wedge (H^A \otimes I_E) \rangle - \int_{M_3} \langle (D^B \Phi_B \otimes I_E) \wedge (K^B \otimes I_E) \rangle. \quad (6)$$

The flexibility in obtaining the Bogomol’nyi energy, a feature not present in the theory of magnetic monopoles, will lead to electric monopoles.

Since topological monopoles, if they exist, are analogous to the BPS magnetic monopole field configurations, we shall use the same symmetry breaking mechanism [2]. We place the solitonic core region at the origin. Let G and H be compact and connected gauge groups, where the group H is assumed to be embedded in G . The gauge group of the core region G is spontaneously broken to H outside of the core region when the Higgs field is covariantly constant, $D\Phi = 0$. In regions far from the core ($r \rightarrow \infty$) where we assume that $D^A \Phi_A = 0$, it can be shown that

$$H^A = \Phi_A F_A, \quad (7)$$

where $F_A \in \Lambda^2(M_3, E_H)$, a two-form on M_3 taking values in the H -Lie algebra bundle, denoted by E_H here [2]. An equivalent expression to (7) can be written when $D^B \Phi_B = 0$. We adopt units so that $\Phi^2 = 1$ when $r \gg 1$ and where spontaneous symmetry breaking has occurred. When $G = U(n)$ and $H = U(1)$, F_A becomes a pure imaginary two-form on M_3 . The Bogomol’nyi solitons defined by (5) have an energy coming from (6) topologically fixed by

$$\begin{aligned} \mathcal{E} &= - \int_{M_3} d \langle \Phi_A H^A \rangle - \int_{M_3} d \langle \Phi_B K^B \rangle \\ &= - \int_{S^2} \langle \Phi_A H^A \rangle - \int_{S^2} \langle \Phi_B K^B \rangle \end{aligned} \quad (8)$$

where S^2 is a large sphere surrounding the monopole core. Details of the solitonic particles in this theory now depend on the extent to which they satisfy the Bogomol’nyi equations (5) and are ‘spontaneously broken’.

Consider first non-singular particle-like solutions to *both* Bogomol'nyi equations in (5). Substituting (7) into (8) and using the normalisation condition $\Phi^2 = 1$ for both Higgs fields, the energy is fixed by $-\int F_A - \int F_B$. Let us conventionally interpret $\int F_A/2\pi$ as the magnetic charge (g), and $\int F_B/2\pi$ as the electric charge (q). We can view F_A and F_B as curvatures on the line bundles determined by Φ_A and Φ_B , respectively, because from (7) they are the projections of H_A and K_B on the line bundles. The magnetic and electric charges are thereby the Chern numbers associated to complex line bundles with curvatures F_A and F_B , respectively, and in this theory are quantized at the classical level. The stability of the solitonic particle is argued from the topological interpretation that can be given to (8). When both equations in (4) are satisfied the topological stability is assured if either the electric or magnetic charge is non-vanishing. To this point, we have used only half of the Bogomol'nyi equations, $D^A\Phi_A = D^B\Phi_B = 0$. From the projective flatness of the curvatures in the Bogomol'nyi equations (5), $H^A = K^B = FI_E$, and (7) we conclude that $F = \varphi_A F_A = \varphi_B F_B$ where $\Phi_A = \varphi_A I_E$ and $\Phi_B = \varphi_B I_E$. From this we find that

$$\int_{S^2} F_A = \int_{S^2} F_B. \quad (9)$$

Therefore non-singular, stable, particle-like solutions to the Bogomol'nyi equations (5) are dyons. To obtain electric monopoles there would appear to be two possibilities, both resulting from a weakening of the Bogomol'nyi equations (5).

In the first possibility, one or the other of the non-abelian gauge fields, H^A or K^B , would not break to $U(1)$ in the far-field, so that non-abelian gauge field would pass unnoticed through the detector. This corresponds to the case where, for example, $D^B\Phi_B = 0$, $H^A = K^B = iFI_E = F_B\Phi_B$, but $D^A\Phi_A \neq 0$. The Bogomol'nyi bound in this case becomes

$$\mathcal{E} = - \int_{S^2} F_B < \Phi_A \Phi_B > - \int_{S^2} F_B. \quad (10)$$

The second term is -2π times the electric charge. Since the projective flatness reduces the gauge group to $U(n)$, we may use the Killing-Cartan form for the bundle inner product, $< >$. The first integral in (10) can be made to vanish when Φ_A is traceless, because $\Phi_B = \varphi_B I_E$. An explicit example of an electric monopole would be $A = B$ projectively flat, $D^B\Phi_B = 0$, and Φ_A is any sufficiently differentiable function taking values in $SU(n)$. The full variational field equations are seen to be satisfied.

The second possibility would allow both gauge fields to break far from the core using $D^A\Phi_A = D^B\Phi_B = 0$, but the two gauge far-fields $H^A = F_A\Phi_A$ and $K^B = F_B\Phi_B$ would be completely decoupled so that F_A and F_B and their respective topological charges would be unrelated. That is, the projective flatness in the Bogomol'nyi equations would be relaxed. The electric and magnetic charges in this case become, respectively,

$$2\pi q = - \int_{S^2} F_B, \quad 2\pi g = - \int_{S^2} F_A,$$

with no relation between q and g analogous to (9).

In this paper we have introduced a topological field theory with a Bogomol'nyi structure permitting BPS electric, magnetic and dyonic monopoles. From the general arguments in [5] the particle spectrum and mass compare favourably with that of the intermediate vector bosons. In this theory $W^\pm$ have energies of opposite sign, so that one is the anti-particle of the other. The Z_0 , presumably, corresponds to the case where $H^A = K^B = iFI_E$, but $D^A\Phi_A \neq 0$ and $D^B\Phi_B \neq 0$. The gauge far-fields for the Z_0 are non-abelian and pass unnoticed through the detectors, and have a stabilizing energy given by

$$\mathcal{E} = - \int_{S^2} F < \Phi_A > - \int_{S^2} F < \Phi_B >$$

We do not yet know whether the classical intermonopole forces share much in common with those for BPS magnetic monopoles, and we do not know whether the classical BPS topological monopole dynamics possess quantum mechanical behaviour as was found for BPS magnetic monopoles in [6].

References

- [1] Atiyah, M., Hitchin, N., The Geometry and Dynamics of Magnetic Monopoles (Princeton U.P., New Jersey, 1988).
- [2] Goddard, P., Olive, D., Magnetic monopoles in gauge theories, *Rep. Prog. Phys.* **41** (1978) pp.1357-1437.
- [3] Horowitz, G.T., Exactly soluble diffeomorphism invariant theories, *Comm. Math. Phys.* **125** (1989) p. 46.

- [4] Kobayashi, S., Differential Geometry of Complex Vector Bundles (Princeton U.P., New Jersey, 1987).
- [5] Montonen, C., Olive, D.I., Magnetic monopoles as gauge particles?, *Phys. Lett.* **72B** (1977) pp.117-120.
- [6] Temple-Raston, M., Alexander, D., Differential cross-sections and escape plots for low-energy SU(2) BPS magnetic monopole dynamics, *Nucl. Phys.* **B397** (1993) pp.195-213.
- [7] Temple-Raston, M., Dyons in topological field theories, *Lett. Math. Phys.* **23** (1991) pp.91-97.
- [8] Temple-Raston, M., Instantons in topological field theories, preprint CON93-3/hep-th:9308055.
- [9] Temple-Raston, M., Solitons in topological field theories, preprint BISH93-10-1/hep-th:9310087.